Contents

Conference Photograph vi

Foreword vii

Conference Information viii

Chapter 1. Quantum Hall effect and related phenomena

Probing the depletion region of a two-dimensional electron system in high magnetic fields
J. Weis, Y.Y. Wei and K. von Klitzing 1

Dynamics of nonequilibrium electrons at the breakdown of the quantum Hall effect
I.I. Kaya, G. Nachtwei, K. von Klitzing and K. Eberl 8

Magnetization of an incompressible two-dimensional electron gas
S.A.J. Wiegers, M. Specht, E.D. Bibow, L.P. Lévy, S. Melinte, E. Grivei, V. Bayot, M.Y. Simmons, D.A. Ritchie, M. Shayegan, A. Cavanna, B. Etienne, G. Martinez and P. Wyder 16

Inhomogeneous transport and derivative relations in the quantum Hall regime
S.H. Simon 23

Magnetoresistance of a two-dimensional electron gas in a one-dimensional superlattice defined by strong modulation fields
S.D.M. Zwerschke and R.R. Gerhardts 28

Angle-resolved ballistic phonon absorption spectroscopy in the lowest Landau level
C.J. Mellor, U. Zeitler, A.M. Devitt, S.H. Roshko, A.J. Kent, K.A. Benedict, T. Cheng and M. Henini 36

Dissipation process and the river formation model of the quantum Hall effect breakdown
C. Chaubet and F. Geniet 43

Quantum Hall effect breakdown: analogies with fluid dynamics
L. Eaves 47

Interactions in the integer quantum Hall effect: Critical conductivity and dynamical scaling
M. Backhaus and B. Huckestein 52

Collapse of quantized Hall resistance at high Hall electric fields
S. Kawaji, H. Iizuka, T. Kuga and T. Okamoto 56

Current distribution in antidot arrays near the breakdown of the quantum Hall effect
J. Gro and R.R. Gerhardts 60

Hierarchical network models for the quantum Hall effect
M. Janssen, R. Merkt, J. Meyer and A. Weymer 65

Conductance noise of submicron wires in the regime of Quantum Hall Effect
J. Wróbel, J. Jaroszyński, T. Dietl, K. Regiński and M. Bugajski 69

High-frequency conductivity experiments on the integer quantum Hall effect at the limits of universality
R. Meisels, F. Kuchar, W. Belitsch, G. Hein and K. Pierz 74

Nonlinear transport properties of samples with inner current contacts in quantum Hall regime
T. Takamasu, Y. Oikawa, G. Kido, A. Endo, M. Kato, S. Katsumoto and Y. Iye 78

New resistance maxima of the 2DEG at fractional filling factors
S. Kronmüller, W. Dietsche, K. von Klitzing, W. Wegscheider and M. Bichler 82

Re-entrant behavior of the $\nu = 4/3$ fractional quantum Hall effect in a front-and-back-gated 2D hole gas
K. Muraki and Y. Hirayama 86

Chapter 2. Spin physics in low dimensional systems

Heat capacity of a 2DEG
E. Grivei, J.-M. Beuken, G. Mariage, V. Bayot and M. Shayegan 90
Electronic properties and quantum Hall effect in multi-layer electron systems
L. Brey 97
Optically detected nuclear magnetic resonance from a single heterojunction in the fractional quantum Hall regime
H.D.M. Davies, R.L. Brockbank, J.F. Ryan and A.J. Turberfield 104
Optically pumped NMR in the quantum Hall regimes
P. Khandelwal, N.N. Kuzma, S.E. Barrett, L.N. Pfeiffer and K.W. West 113
Ultraslow electron spin dynamics in the fractional quantum hall regime
N.N. Kuzma, P. Khandelwal, S.E. Barrett, L.N. Pfeiffer and K.W. West 121
Quantum transport in spin-singlet $\nu = 2/3$ edges
K. Imura and N. Nagaosa 125
A study of the relative strengths of spin-pseudospin phases in a strongly coupled double quantum well system
C.H.W. Barnes, A.G. Davies, K.R. Zolleis, M.Y. Simmons and D.A. Ritchie 130
Optical measurements of spin polarization in a 2D electron system with g-factor tunable through zero
V. Huard, S. Lovisa, R.T. Cox, K. Saminadayar, M. Potemski, A. Arnoult, J. Cibert, S. Tatarenko and A. Wasiela 136

Chapter 3. Quantum dots

Transport spectroscopy of lateral few electron quantum dots in a magnetic field
C. Gould, P. Hawrylak, A.S. Sachrajda, Y. Feng and Z. Wasilewski 141
Shell structure and exchange-related spin effects for quantum dots in a magnetic field
O. Steffens, M. Suhrke and U. Rössler 147
Quantum dots with interacting electrons: Energy spectra and magnetization
W. Sheng and H. Xu 152
Spin states in circular and elliptical quantum dots
S. Sasaki, D.G. Austing and S. Tarucha 157
Coalescence of extended states in large quantum dots
V. Kagalovsky, B. Horovitz and Y. Avishai 161
AC Kondo effect in quantum dots
R. Aguado, R. López, G. Platero and C. Tejedor 165
Magnetic field effects in InP self-assembled quantum dots
M. Sugisaki, H.-W. Ren, K. Nishi, S. Sugou, T. Okuno and Y. Masumoto 169
Singlet–triplet transitions in a few-electron quantum dot
W.G. van der Wiel, T.H. Oosterkamp, J.W. Janssen, L.P. Kouwenhoven, D.G. Austing, T. Honda and S. Tarucha 173
Magneto-luminescence of quasi-zero dimensional $In_{0.25}Ga_{0.75}As/GaAs$ quantum dots
H.H. Cheng, R.J. Nicholas, A. Priest, F.Y. Tsai, C.P. Lee and J. Sanchez-Dehesa 178
A quantum dot in the limit of strong coupling to reservoirs
J. Schmid, J. Weis, K. Eberl and K. von Klitzing 182
Stability of interacting electrons in a magnetic quantum dot
G.P. Mallon and P.A. Maksym 186
Electronic structure of self-assembled quantum dots in high magnetic fields
A. García-Cristóbal, V.M. Fomin and J.T. Devreese 190

Spin blockade in quantum dots in magnetic fields
H. Imamura, H. Aoki and P.A. Maksym 194
Magneto-thermopower of a chaotic, ballistic quantum dot
H. Buhmann, S. Möller, S.F. Godijn and L.W. Molenkamp 198
Photoluminescence measurements of InP quantum dots in pulsed magnetic fields
R. Provoost, M. Hayne, M.K. Zundel, K. Eberl and V.V. Moshchalkov 203

Chapter 4. Transport properties of low dimensional systems

Minigaps and the quantum Hall effect in broken gap InAs/GaSb heterostructures
R.J. Nicholas, M. Lakrimi, S.W. Khym, N.J. Mason, A.J.L. Poulter, T. Vaughan, P.J. Walker, D.K. Maude, J.C. Portal, D.M. Symons and F.M. Peeters 207
Surface acoustic wave wavelength and temperature dependence of the induced periodic density modulation effect at $\nu = 1/2$
R.L. Willett, L.N. Pfeiffer and K.W. West 215
Coulomb drag in a quasiclassical magnetic field
A.V. Khaetskii and Yu.V. Nazarov 220
Magnetic Breakdown in the semimetallic InAs/GaSb system
D.M. Symons, M. Lakrimi, R.J. Nicholas, D.K. Maude, J.C. Portal, N.J. Mason and P.J. Walker 225
Diamagnetic effects in structures containing a single, a few or an infinite number of coupled delta layers
P.M. Koenraad, A.B. Henriques, A.F.W. van de Stadt and J.H. Wolter 229
Magnetic field induced charge instabilities in weakly coupled superlattices
R. Aguado and G. Platero 233
Landau and spin levels in InAs quantum wells resolved with in-plane and parallel magnetic fields
S. Brosig, K. Ensslin, B. Brar, M. Thomas and H. Kroemer 239
Magnetotransport in GaAs δ-doped by Sn
R.T.F. van Schaijk, A. de Visser, V.A. Kulbachinskii, V.G. Kytin, R.A. Lunin, V.G. Mokerov, A.S. Bugaev and A.P. Senichkin 243
Magnetooscillations in a wide parabolic well in the presence of an in-plane magnetic field
G.M. Gusev, J.R. Leite, N.T. Moshegov and A.I. Toropov 248
Investigation of the spatial variation of scattering centers in parabolic quantum wells
T. Heinzel, G. Salis, P. Wirth, K. Ensslin, K. Maranowski and A.C. Gossard 252
Optical probing of the minigap in InAs/GaSb superlattices
A.J.L. Poulter, M. Lakrimi, R.J. Nicholas, N.J. Mason and P.J. Walker 256
Anomalous coincidences between valley split Landau levels in a Si/SiGe heterostructure
H.W. Schumacher, A. Nauen, U. Zeitler, R.J. Haug, P. Weitz, A.G.M. Jansen and F. Schäffler 260
Perpendicular and parallel magnetic field transport measurements in gated InAs/GaSb DHETs
M. Lakrimi, J. Rehman, D.M. Symons, R.J. Nicholas, N.J. Mason and P.J. Walker 264
Magneto-photoconductivity of a 2DEG under intense terahertz radiation
P.M. Koenraad, R.A. Lewis, L.R.C. Waumans, C.J.G.M. Langerak, W. Xu and J.H. Wolter 268
Anisotropic magneto-resistance of laterally modulated GaAs/AlGaAs systems with a 15–20 nm periodicity formed on vicinal (1 1 1)B substrates
Y. Nakamura and H. Sakaki 273
Quantization and chiral edge state properties in nearly 3D quantum well structures
B. Zhang, J.S. Brooks, Z. Wang, J.A.A.J. Perenboom, J. Simmons, J. Reno, N. Lumpkin, J. O'Brien and R. Clark 279

Chapter 5. Optical properties of low dimensional systems

Magneto-optics of a two-dimensional electron gas
M. Potemski 283

High-field cyclotron resonance and electron–phonon interaction in modulation-doped multiple quantum well structures
Y.J. Wang, Z.X. Jiang, B.D. McCombe, F.M. Peeters, X.G. Wu, G.Q. Hai, T.J. Eustis and W. Schaff 292
Spatially resolved magneto-optics on confined systems
A. Zrenner, M. Markmann, A. Paassen, A.L. Efros, M. Bichler, W. Wegscheider, G. Böhm and G. Abstreiter 300
Magneto-optical study of excitons in quantum wells, wires and dots in high magnetic fields
N. Miura, H. Kunimatsu, K. Uchida, Y. Matsuda, T. Yasuhira, H. Nakashima, Y. Sakuma, Y. Awano, T. Futatsugi and N. Yokoyama 308
Photoluminescence in asymmetric quantum wells at $\nu > 1$
K. Asano and T. Ando 319
Magneto-optical study of ZnSe-based quantum wells
W. Ossau, D.R. Yakovlev, U. Zehnder, G.V. Astakhov, A.V. Platonov, V.P. Kochereshko, J. Nürnberger, W. Faschinger, M. Keim, A. Waag, G. Landwehr, P.C.M. Christianen, J.C. Maan, N.A. Gippius and S.G. Tikhodeev 323
Excitonic and free-carrier recombination of a two-dimensional electron gas in high magnetic fields
M. Hayne, C.L. Jones, R. Bogaerts, A. Usher, F. Herlach, V.V. Moshchalkov and M. Henini 327
Magnetic field effects on the Rabi splitting and radiative decay rates of the exciton–polariton states in a semiconductor microcavity
H. Fenniche, S. Jaziri and R. Bennaceur 331
Landau quantized 2D nonlocal Bernstein mode magnetoplasmon coupling with surface and bulk phonons
N.J.M. Horing and Godfrey Gumbs 335
Experiments on semiconductor systems using a pulsed-field-magnet free-electron-laser combination
C.J.G.M. Langerak, J. Singleton, L. Li, L. Van Bockstal, A. Ardavan, M.J. van der Pol, A.F.G. van der Meer and F. Herlach 339
Angular dependent cyclotron resonance of two-dimensional electron gas in GaAs/AlGaAs quantum wells in high magnetic fields up to 150 T
H. Arimoto, T. Saku, Y. Hirayama and N. Miura 343
Magneto-photoluminescence of GaAs/AlGaAs heterojunctions under hydrostatic pressure
A.N. Priest, R.J. Nicholas, D. Symons, J.C. Portal, C.T. Foxon and J.J. Harris 347
Confined photonic modes in distributed Bragg microresonator in magnetic field
N.A. Gippius and S.G. Tikhodeev 351
Cyclotron-stark resonances in superlattices under the influence of strong THz radiation
P. Kleinert and V.V. Bryksin 356
Pressure dependence of the cyclotron resonance modes in high mobility two-dimensional hole systems in GaAs–(Ga,Al)As heterojunctions
G.L.B. Verschoor, J.M. Schrama, A.V. Semeno, A. Kornilov, A-K. Klehe, A. Ardavan, J. Singleton, P. Goy, J.M. Chamberlain, M. Henini and T. Cheng 359
Photoluminescence from $Si/Si_{1-x}Ge_x$ single quantum wells in high magnetic fields: localized and free exciton recombination
C. Penn, P.C.M. Christianen, F. Schäffler, J.C. Maan and G. Bauer 363
Magneto-optical oscillations in the photoluminescence of quantum wells and the resonant polaron effect
A.R. Alves, L.A. Cury, P.S.S. Guimarães and M.V.B. Moreira 367

Chapter 6. Mesoscopic systems and phenomena

Coherent properties of in plane gated submicron rings in a small-signal regime
A.A. Bykov, I.V. Marchishin, A.G. Pogosov, L.V. Litvin, E.B. Olshanetskii and G.M. Gusev 371
Quantum analog of channeled electron trajectories in periodic magnetic and electric fields
A. Manolescu, S.D.M. Zwerschke, M. Niţă, U.J. Gossmann and R.R. Gerhardts 375
Induced magnetoresistance in semiconductor devices due to single sub-micron magnetic barriers
V. Kubrak, F. Rahman, N. Overend, B.L. Gallagher, P.C. Main, J. Boeck, M. Behest, C.H. Marrows and M.A. Howson 380

Coupled one-dimensional subbands in in-plane magnetic fields
G. Salis, T. Heinzel, K. Ensslin, O. Homan, W. Bächtold, K. Maranowski and A.C. Gossard 384
Aharonov–Bohm oscillation and periodic orbits in antidot lattices
S. Uryu and T. Ando 388
Longitudinal magnetoresistance in magnetic barrier systems
K. Tsubaki 392
Antidot shape dependence of the commensurability oscillation of magnetoresistance in two-dimensional antidot arrays
T. Azuma and T. Osada 397
Universal conductance fluctuations in a parabolic quantum well tunable from two- to three-dimensional behavior
T. Ihn, M. Huberty, G. Salis, K. Ensslin, K. Maranowski and A.C. Gossard 401
Observation of a new low-field minimum in magnetically modulated systems
J. Raabe, D. Weiss and V. Umanski 405
Magnetotransport in large diameter InAs/GaSb antidot lattices
J. Eroms, M. Zitzlsperger, D. Weiss, J.H. Smet, C. Albrecht, R. Fleischmann, M. Behet, J. De Boeck and G. Borghs 409
Phase coherence in double quantum well mesoscopic wires
S.T. Stoddart, P.C. Main, M.J. Gompertz, A. Nogaret, L. Eaves, M. Henini and S.P. Beaumont 413

Chapter 7. Many-body phenomena

Probing the transition from insulating to metallic behaviour using bi-layer electron systems
A.G. Davies, C.H.W. Barnes, K.R. Zolleis, M.Y. Simmons and D.A. Ritchie 417
Many-particle ground states and excitations in nanometer-size quantum structures
A. Lorke and R.J. Luyken 424
Electron correlations in quasi-2D structures at high magnetic fields in the presence of embedded donor impurities
Z.X. Jiang, B.D. McCombe and P. Hawrylak 431
Kinetics and localization of composite fermions in a random magnetic field
D.G. Polyakov, F. Evers, A.D. Mirlin and P. Wölfle 441
High magnetic field far-infrared spectroscopy of spatially separated electron–hole layers
Yu.B. Vasilyev, S.D. Suchalkin, K. von Klitzing, B.Ya. Meltser, S.V. Ivanov and P.S. Kop'ev 445
Asymmetry of elementary interactions in 2D systems in a magnetic field
O.V. Kibis 449
Edge magnetoplasmons in double quantum wells: Influence of Coulomb coupling and tunneling
P. Vasilopoulos and O.G. Balev 452
Cyclotron resonance in high mobility CdTe/CdMgTe 2D electron system in integer quantum Hall regime
Y. Imanaka, T. Takamasu, G. Kido, G. Karczewski, T. Wojtowicz and J. Kossut 457
Many-body effects in photoluminescence of a two-dimensional electron gas in high magnetic field
H.P. van der Meulen, J. Rubio, I. Azcona, J.M. Calleja, P.C.M. Christianen and J.C. Maan 462
Dynamics of charged excitons in GaAs quantum wells under high magnetic field
V. Ciulin, G. Finkelstein, S. Haacke, J.-D. Ganière, V. Umansky, I. Bar-Joseph and B. Deveaud 466
Inhibited recombination of negatively-charged excitons in GaAs quantum wells at high magnetic fields
H. Okamura, D. Heiman, M. Sundaram and A.C. Gossard 470
Collective excitations in a 2D electron system: Canted field geometry
D.C. Marinescu, J.J. Quinn and A. Wójs 474
Multiply-charged magnetoexcitons in quantum-well systems
A.V. Chaplik and A.O. Govorov 477
Far-infrared studies of extremely high mobility gated GaAs/AlGaAs structures in magnetic fields
R.J. Heron, R.A. Lewis, R.G. Clark, R.P. Starrett, B.E. Kane, G.R. Facer, N.E. Lumpkin, D.G. Rickel, L.N. Pfeiffer and K.W. West 481

Cyclotron masses of asymmetrically doped HgTe quantum wells
A. Pfeuffer-Jeschke, F. Goschenhofer, S.J. Cheng, V. Latussek, J. Gerschütz, C.R. Becker, R.R. Gerhardts and G. Landwehr 486
Incompressible states of negatively charged magneto-excitons
A. Wójs, P. Hawrylak and J.J. Quinn 490
Coupled cyclotron resonance transitions of bilayer 2DEG systems in GaAs
C.M. Hu, E. Batke, S.C. Shen, K. Köhler and P. Ganser 494
Instability of the QHE induced by a nearby two-dimensional-electron system
C. Jörger, W. Wegscheider, W. Dietsche and K. von Klitzing 499
Exchange interaction in degenerate p-type quantum wells
M. Kemerink, P.M. Koenraad and J.H. Wolter 503

Chapter 8. Tunneling phenomena

Magneto-tunnelling spectroscopy of a two-dimensional electron system
P.C. Main, A.S.G. Thornton, T. Ihn, L. Eaves, K.A. Benedict and M. Henini 507
Limitation of tunneling between edge states by a mesoscopic island in narrow quantum Hall conductors
Z.H. Liu, G. Nachtwei, J. Gro, R.R. Gerhardts, J. Weis, K. von Klitzing and K. Eberl 514
Fermi edge singularities in high magnetic fields
K.A. Benedict, A.S.G. Thornton, T. Ihn, P.C. Main, L. Eaves and M. Henini 519
Magnetic field dependent zero-bias diffusive anomaly in Pb-oxide–*n*-InAs structures: Coexistence of 2D and 3D states
G.M. Minkov, A.V. Germanenko, S.A. Negachev, O.E. Rut and E.V. Sukhorukov 523
Resonant magnetic field induced enhancement of the tunneling current in multi-quantum wells
G.S. Vieira, P.S.S. Guimarães, E.S. Alves, S.J. Allen, K.L. Campman and A.C. Gossard 527
Tunneling current bistability of correlated 2D electron–hole layers
A. Parlangeli, P.C.M. Christianen, A.K. Geim, J.C. Maan, L. Eaves, P.C. Main and M. Henini 531
Spin dependence of the interlayer tunneling in double quantum wells in the quantum Hall regime
S. Kishimoto, Y. Ohno, F. Matsukura and H. Ohno 535
Polaron pinning effects in superlattices at high electric and magnetic fields
R.K. Hayden, A. Nogaret, L. Eaves, N. Miura and M. Henini 540
A study of miniband conduction in Wannier–Stark superlattices at high magnetic fields
H.M. Murphy, A. Nogaret, S.T. Stoddart, L. Eaves, P.C. Main, M. Henini, D.K. Maude, N. Mori, J.-C. Portal and C. Hamaguchi 544

Chapter 9. Semimagnetic semiconductors

Magnetic structure of short-period CdTe/MnTe superlattices
H. Krenn, R. Rupprecht, S. Holl, W. Faschinger, H. Pascher and G. Bauer 548
Magnetic field mediated anti-Stokes luminescence of semimagnetic ADQW structures
W. Heimbrodt and M. Happ 553
Magnetooptics of CdTe- and (Cd,Mn)Te- based modulation doped quantum well structures
V.P. Kochereshko, A.V. Platonov, D.R. Yakovlev, T. Wojtowicz, M. Kutrowski, G. Karczewski, J. Kossut, W. Ossau and G. Landwehr 557
Magnetotunneling spectroscopy of resonant tunneling diode using ferromagnetic (Ga,Mn)As
N. Akiba, F. Matsukura, Y. Ohno, A. Shen, K. Ohtani, T. Sakon, M. Motokawa and H. Ohno 561
Cyclotron resonance in $Cd_{1-x}Fe_xS$ and $Ga_{1-x}Mn_xAs$ at megagauss magnetic fields
Y.H. Matsuda, H. Arimoto, N. Miura, A. Twardowski, H. Ohno, A. Shen and F. Matsukura 565
ESR study of Mn doped II–VI and III–V DMS
H. Nojiri, M. Motokawa, S. Takeyama, F. Matsukura and H. Ohno 569
Magnetotransport properties of all semiconductor (Ga,Mn)As/(Al,Ga)As/(Ga,Mn)As tri-layer structures
F. Matsukura, N. Akiba, A. Shen, Y. Ohno, A. Oiwa, S. Katsumoto, Y. Iye and H. Ohno 573
Magneto-optics of a two-dimensional electron gas confined in $Cd_{1-x}Mn_xTe$ quantum wells
F.J. Teran, M.L. Sadowski, M. Potemski, G. Karczewski, S. Mackowski and J. Jaroszynski 577

Chapter 10. Metal–insulator transition

Magnetic field suppression of the conducting state in two dimensions
S.V. Kravchenko, D. Simonian, M.P. Sarachik, A.D. Kent and V.M. Pudalov 582
Hall effect of pinned and depinned 2-D electron and hole solids
F. Perruchot, C.J. Mellor, R. Gaál, B. Sas, G. Deville, B. Etienne, B.L. Gallagher, M. Henini, C.T. Foxon, J.J. Harris and F.I.B. Williams 587
Quantum-Hall to insulator transition
H. Potempa, A. Bäker and L. Schweitzer 591
2D lattice of coupled Sinai billiards: metal or insulator at g 1?
M.V. Budantsev, Z.D. Kvon, A.G. Pogosov, G.M. Gusev, J.C. Portal, D.K. Maude, N.T. Moshegov and A.I. Toropov 595
Magnetic field induced metal-to-insulator transition in wide quantum wells
P. Ganitzer, A. Homer, M. Lucyshyn, B. Jamnig, J. Oswald, D.K. Maude and J.C. Portal 600
On the extended states in the quantum Hall regime and in zero magnetic field
H. Nielsen 604
Magnetoconductance of the anomalous 2D conducting phase in parallel field
D. Simonian, S.V. Kravchenko, K.M. Mertes, M.P. Sarachik and V.M. Pudalov 607

Chapter 11. Novel materials, bulk semiconductors, etc.

Vortex matter in mesoscopic superconductors
J.J. Palacios 610
Magnetic levitation experiments in Tohoku University
M. Motokawa, I. Mogi, M. Tagami, M. Hamai, K. Watanabe and S. Awaji 618
Field-induced reentrant electronic phase transitions in graphite
H. Yaguchi and J. Singleton 621
Effective mass and bandstructure of n-InAs from magnetophonon resonance and Raman scattering at temperatures between $T = 64$ and 360 K
D. Schneider, C. Brink, G. Irmer and P. Verma 625
Band structure of β-HgS from Shubnikov–de Haas effect
K. Dybko, W. Szuszkiewicz, E. Dynowska, W. Paszkowicz and B. Witkowska 629
Semiclassical angular effect of magnetoresistance and field-induced confinement state in the organic quasi-one-dimensional conductors
T. Osada 633
Spin resonance studies of the quasi-one-dimensional Heisenberg antiferromagnet Cs_2CuCl_4
J.M. Schrama, A. Ardavan, A.V. Semeno, P.J. Gee, E. Rzepniewski, J. Suto, R. Coldea, J. Singleton and P. Goy 637
Magnetic versus hyperfine field in the anomalous Hall effect
H. Bednarski, V. Fleurov and I.D. Vagner 641
Magneto-plasmon spectrum in a semiconductor under intense laser radiation
W. Xu, C. Zhang and R.A. Lewis 645
A new kind of magnetic resonance observed in the organic molecular metal -$(BEDT\text{-}TTF)_2KHg(SCN)_4$
A. Ardavan, J.M. Schrama, A. Semeno, S.J. Blundell, J. Singleton, P. Goy, M. Kurmoo and P. Day 649
Spontaneous current oscillations in n-GaAs in high magnetic fields
V.A. Samuilov, G. Kiss, V.K. Ksenevich, G. Remenyi and B. Podor 654
Magneto-optical investigations on cubic GaN in high magnetic fields
I. Stolpe, N. Puhlmann, H.-U. Müller, O. Portugall, M. von Ortenberg, D. Schikora, D.J. As, B. Schöttker and R. Lischka 659

List of Contributors 663

Subject Index 667